

cse371/mat371
LOGIC

Professor Anita Wasilewska

Fall 2017

LECTURE 5a

Chapter 5
HILBERT PROOF SYSTEMS: Completeness of Classical
Propositional Logic

Lecture 5a

PART 1: Introduction

PART 2: **Proof** of the **Main Lemma**

PART 3: **Proof 1**: Constructive Proof of **Completeness
Theorem**

PART 1: Introduction

Two Proofs

There are **many proof systems** that describe classical propositional logic, i.e. that are **complete proof systems** with the respect to the classical semantics.

We present here a **Hilbert proof system** for the classical propositional logic and discuss **two ways** of proving the **Completeness Theorem** for it.

Any **proof** of the **Completeness Theorem** consists always of **two parts**.

Two Proofs

First we have show that **all formulas that have a proof are tautologies.**

This implication is also called a **Soundness Theorem**, or **Soundness Part** of the **Completeness Theorem**

The second implication says: **if a formula is a tautology then it has a proof.**

This alone is sometimes called a **Completeness Theorem** (on assumption that the system is sound)

Traditionally it is called a **completeness part** of the **Completeness Theorem**

Two Proofs

The **proof** of the soundness part is standard.

We concentrate here on the **completeness part** of the **Completeness Theorem** and present **two proofs** of it

The **first proof** is straightforward. It shows how one can use the assumption that a formula **A is a tautology** in order to **construct** its **formal proof**

It is hence called a **proof - construction method**.

Two Proofs

The **second proof** shows how one can **prove** that a formula **A is not a tautology** **from** the fact that **it does not have a proof**

It is hence called a **counter-model construction method**.

All these **proofs** and considerations are **relative** to proof systems and their **semantics**

At this moment the semantics is **classical** and the proof system is **H_2**

Reminder: we write $\models A$ to denote that **A** is a **classical tautology**

Proof System H_2

Reminder: H_2 is the following proof system:

$$H_2 = (\mathcal{L}_{\{\Rightarrow, \neg\}}, \mathcal{F}, \{A1, A2, A3\}, MP)$$

The axioms **A1 – A3** are defined as follows.

$$\mathbf{A1} \quad (A \Rightarrow (B \Rightarrow A)),$$

$$\mathbf{A2} \quad ((A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))),$$

$$\mathbf{A3} \quad ((\neg B \Rightarrow \neg A) \Rightarrow ((\neg B \Rightarrow A) \Rightarrow B)))$$

$$(MP) \quad \frac{A ; (A \Rightarrow B)}{B}$$

Proof System H_2

Obviously, the selected axioms A_1, A_2, A_3 are **tautologies**, and the **MP** rule leads from tautologies to tautologies.

Hence our proof system H_2 is **sound** and the following theorem holds.

Soundness Theorem

For every formula $A \in \mathcal{F}$,

If $\vdash_{H_2} A$, then $\models A$

System H_2 LEMMA

We have proved in **Lecture 5** the following

Lemma

The following formulas **are provable** in H_2

1. $(A \Rightarrow A)$
2. $(\neg\neg B \Rightarrow B)$
3. $(B \Rightarrow \neg\neg B)$
4. $(\neg A \Rightarrow (A \Rightarrow B))$
5. $((\neg B \Rightarrow \neg A) \Rightarrow (A \Rightarrow B))$
6. $((A \Rightarrow B) \Rightarrow (\neg B \Rightarrow \neg A))$
7. $(A \Rightarrow (\neg B \Rightarrow (\neg(A \Rightarrow B))))$
8. $((A \Rightarrow B) \Rightarrow ((\neg A \Rightarrow B) \Rightarrow B))$
9. $((\neg A \Rightarrow A) \Rightarrow A)$

First Proof

The **first proof** of **Completeness Theorem** presented here is very **elegant** and **simple**, but is **applicable only** to the **classical propositional logic**

This proof is, as was the proof of **Deduction Theorem**, a fully **constructive**

The technique it **uses** , because of its specifics **can't be used** even in a case of classical **predicate logic**, not to mention variety of **non-classical** logics

Second Proof

The **second proof** is much more complicated.

Its **strength** and **importance** lies in a fact that the methods it uses can be applied in an extended version to the **proof of completeness** for **classical predicate logic** and some **non-classical** propositional and predicate logics

The way **we define** a **counter-model** for any **non-provable A** is general and non-constructive

We call it a **counter-model existence method**

PART 2: Proof of the MAIN LEMMA

Completeness Theorem

The **proof** of the **Completeness Theorem** presented here is similar in its structure to the proof of the **Deduction Theorem** and is due to **Kalmar, 1935**

It is a **constructive proof**

It **shows** how one can use the assumption that a formula **A** is **a tautology** in order to **construct** its formal proof.

We hence call it a **proof construction method**. It relies heavily on the **Deduction Theorem**

It is **possible** to prove the **Completeness Theorem** **independently** from the **Deduction Theorem** and we will present two of such a proofs in later chapters.

Introduction

We first present **one definition** and prove **one lemma**

We write $\vdash A$ instead of $\vdash_S A$ as the system S is fixed.

Let A be a formula and b_1, b_2, \dots, b_n be all propositional variables that occur in A , i.e.

$$A = A(b_1, b_2, \dots, b_n)$$

MAIN LEMMA: Definition 1

Definition 1

Let v be a truth assignment $v : VAR \rightarrow \{T, F\}$

We define, for A, b_1, b_2, \dots, b_n and truth assignment v corresponding formulas A', B_1, B_2, \dots, B_n as follows:

$$A' = \begin{cases} A & \text{if } v^*(A) = T \\ \neg A & \text{if } v^*(A) = F \end{cases}$$

$$B_i = \begin{cases} b_i & \text{if } v(b_i) = T \\ \neg b_i & \text{if } v(b_i) = F \end{cases}$$

for $i = 1, 2, \dots, n$

Example 1

Let A be a formula $(a \Rightarrow \neg b)$

Let v be such that $v(a) = T, v(b) = F$

In this case we have that $b_1 = a, b_2 = b$, and

$$v^*(A) = v^*(a \Rightarrow \neg b) = v(a) \Rightarrow \neg v(b) = T \Rightarrow \neg F = T$$

The corresponding A', B_1, B_2 are:

$$A' = A \quad \text{as } v^*(A) = T$$

$$B_1 = a \quad \text{as } v(a) = T$$

$$B_2 = \neg b \quad \text{as } v(b) = F$$

Example 2

Let A be a formula $((\neg a \Rightarrow \neg b) \Rightarrow c)$

and let v be such that $v(a)=T, v(b)=F, v(c)=F$

Evaluate A', B_1, \dots, B_n as defined by the **definition 1**

In this case $n = 3$ and $b_1 = a, b_2 = b, b_3 = c$

and we evaluate

$$v^*(A) = v^*((\neg a \Rightarrow \neg b) \Rightarrow c) = ((\neg v(a) \Rightarrow \neg v(b)) \Rightarrow v(c)) = ((\neg T \Rightarrow \neg F) \Rightarrow F) = (T \Rightarrow F) = F$$

The corresponding A', B_1, B_2, B_3 are:

$$A' = \neg A = \neg((\neg a \Rightarrow \neg b) \Rightarrow c) \text{ as } v^*(A) = F$$

$$B_1 = a \text{ as } v(a) = T, B_2 = \neg b \text{ as } v(b) = F, \text{ and}$$

$$B_3 = \neg c \text{ as } v(c) = F$$

MAIN LEMMA

The **lemma** stated below **describes a method** of transforming a **semantic notion** of a **tautology** into a **syntactic notion** of **provability**

It **defines**, for any formula A and a truth assignment v a corresponding **deducibility relation**

Main Lemma

For any formula $A = A(b_1, b_2, \dots, b_n)$ and any truth assignment v

If A', B_1, B_2, \dots, B_n are corresponding formulas defined by **definition 1**, then

$$B_1, B_2, \dots, B_n \vdash A'$$

Examples

Example 3

Let A, v be as defined in the **Example 1**, i.e. $A' = A$,
 $B_1 = a$, $B_2 = \neg b$

Main Lemma asserts that

$$a, \neg b \vdash (a \Rightarrow \neg b)$$

Example 4

Let A, v be defined as in **Example 2**, then the **Main Lemma** asserts that

$$a, \neg b, \neg c \vdash \neg((\neg a \Rightarrow \neg b) \Rightarrow c)$$

Proof of the Main Lemma

The proof is by **induction** on the **degree of the formula** A

Base Case $n = 0$

In this case A is **atomic** and so consists of a single propositional variable, say a

If $v^*(A) = T$ then we have by **definition 1**

$$A' = A = a, B_1 = a$$

We obtain, by **definition of provability** from a set Γ of hypothesis for $\Gamma = \{a\}$ that

$$a \vdash a$$

Proof of the Main Lemma

If $v^*(A) = F$ we have by **Definition 1** that

$$A' = \neg A = \neg a \quad \text{and} \quad B_1 = \neg a$$

We obtain, by **definition of provability** from a set Γ of hypothesis for $\Gamma = \{\neg a\}$ that

$$\neg a \vdash \neg a$$

This **proves** that **Main Lemma** holds for $n=0$

Proof of the Main Lemma

Inductive Step

Now **assume** that the **Lemma** holds for **any formula** with $j < n$ connectives

Need to prove: the **Lemma** holds for **A** with n connectives

There are several sub-cases to deal with

Case: **A** is $\neg A_1$

By the **inductive assumption** we have the formulas

$$A'_1, B_1, B_2, \dots, B_n$$

corresponding to the A_1 and the propositional variables b_1, b_2, \dots, b_n in A_1 , such that

$$B_1, B_2, \dots, B_n \vdash A'_1$$

Observe that the formulas **A** and $\neg A_1$ have the same **propositional variables**

So the **corresponding** formulas B_1, B_2, \dots, B_n are **the same** for both of them.

Proof of the Main Lemma

We are going to show that the **inductive assumption** allows us to prove that

$$B_1, B_2, \dots, B_n \vdash A'$$

There are **two cases** to consider.

Case: $v^*(A_1) = T$

If $v^*(A_1) = T$ then by **definition 1** $A'_1 = A_1$ and by the **inductive assumption**

$$B_1, B_2, \dots, B_n \vdash A_1$$

In this case: $v^*(A) = v^*(\neg A_1) = \neg v^*(T) = F$

So we have that $A' = \neg A = \neg\neg A_1$

Proof of the Main Lemma

By Lemma 5. we have that that

$$\vdash (A_1 \Rightarrow \neg\neg A_1)$$

we obtain by the **monotonicity** that also

$$B_1, B_2, \dots, B_n \vdash (A_1 \Rightarrow \neg\neg A_1)$$

By **inductive assumption** $B_1, B_2, \dots, B_n \vdash A_1$ and by **MP** we have

$$B_1, B_2, \dots, B_n \vdash \neg\neg A_1$$

and as $A' = \neg A = \neg\neg A_1$ we get

$$B_1, B_2, \dots, B_n \vdash \neg A \quad \text{and so} \quad B_1, B_2, \dots, B_n \vdash A'$$

Proof of the Main Lemma

Case: $v^*(A_1) = F$

If $v^*(A_1) = F$ then $A'_1 = \neg A_1$ and $v^*(A) = T$ so $A' = A$

Therefore by the **inductive assumption** we have that

$$B_1, B_2, \dots, B_n \vdash \neg A_1$$

that is as $A = \neg A_1$

$$B_1, B_2, \dots, B_n \vdash A'$$

Proof of the Main Lemma

Case: A is $(A_1 \Rightarrow A_2)$

If A is $(A_1 \Rightarrow A_2)$ then A_1 and A_2 have less than n connectives

$A = A(b_1, \dots, b_n)$ so there are some **subsequences** c_1, \dots, c_k and d_1, \dots, d_m for $k, m \leq n$ of the sequence b_1, \dots, b_n such that

$$A_1 = A_1(c_1, \dots, c_k) \quad \text{and} \quad A_2 = A_2(d_1, \dots, d_m)$$

Proof of the Main Lemma

A_1 and A_2 have less than n connectives and so by the **inductive assumption** we have appropriate formulas C_1, \dots, C_k and D_1, \dots, D_m such that

$$C_1, C_2, \dots, C_k \vdash A_1' \quad \text{and} \quad D_1, D_2, \dots, D_m \vdash A_2'$$

and $C_1, C_2, \dots, C_k, D_1, D_2, \dots, D_m$ are **subsequences** of formulas B_1, B_2, \dots, B_n corresponding to the propositional variables in A

By **monotonicity** we have the also

$$B_1, B_2, \dots, B_n \vdash A_1' \quad \text{and} \quad B_1, B_2, \dots, B_n \vdash A_2'$$

Now we have the following sub-cases to consider

Proof of the Main Lemma

Case: $v^*(A_1) = v^*(A_2) = T$

If $v^*(A_1) = T$ then $A_1' = A_1$ and

if $v^*(A_2) = T$ then $A_2' = A_2$

We also have $v^*(A_1 \Rightarrow A_2) = T$ and so $A' = (A_1 \Rightarrow A_2)$

By the above and the **inductive assumption**

$$B_1, B_2, \dots, B_n \vdash A_2$$

and since we have assumed **1.** about **S** and by **monotonicity** we have

$$B_1, B_2, \dots, B_n \vdash (A_2 \Rightarrow (A_1 \Rightarrow A_2))$$

By above and **MP** we have $B_1, B_2, \dots, B_n \vdash (A_1 \Rightarrow A_2)$
that is

$$B_1, B_2, \dots, B_n \vdash A'$$

Proof of the Main Lemma

Case: $v^*(A_1) = T$, $v^*(A_2) = F$

If $v^*(A_1) = T$ then $A_1' = A_1$ and

if $v^*(A_2) = F$ then $A_2' = \neg A_2$

Also we have in this case $v^*(A_1 \Rightarrow A_2) = F$ and so

$A' = \neg(A_1 \Rightarrow A_2)$

By the **above**, the **inductive assumption** and **monotonicity**

$B_1, B_2, \dots, B_n \vdash \neg A_2$

By Lemma 7. and by **monotonicity** we have

$$B_1, B_2, \dots, B_n \vdash (A_1 \Rightarrow (\neg A_2 \Rightarrow \neg(A_1 \Rightarrow A_2)))$$

By above and **MP twice** we have

$B_1, B_2, \dots, B_n \vdash \neg(A_1 \Rightarrow A_2)$ that is

$$B_1, B_2, \dots, B_n \vdash A'$$

Proof of the Main Lemma

Case: $v^*(A_1) = F$

Observe that if $v^*(A_1) = F$ then A_1' is $\neg A_1$ and, whatever value v gives A_2 , we have

$$v^*(A_1 \Rightarrow A_2) = T$$

So A_1' is $(A_1 \Rightarrow A_2)$

Therefore

$$B_1, B_2, \dots, B_n \vdash \neg A_1$$

From Lemma 6. and by monotonicity we have

$$B_1, B_2, \dots, B_n \vdash (\neg A_1 \Rightarrow (A_1 \Rightarrow A_2))$$

Proof of the Main Lemma

By Modus Ponens we get that

$$B_1, B_2, \dots, B_n \vdash (A_1 \Rightarrow A_2)$$

that is

$$B_1, B_2, \dots, B_n \vdash A'$$

We have covered **all cases** and, by **mathematical induction** on the degree of the formula **A** we got

$$B_1, B_2, \dots, B_n \vdash A'$$

The **proof** of the **Main Lemma** is complete

PART3

Proof 1: Constructive Proof of Completeness Theorem

Proof of Completeness Theorem

Now we use the **Main Lemma** to prove the **Completeness Theorem** i.e. to prove the following implication

For any formula $A \in \mathcal{F}$

if $\models A$ then $\vdash A$

Proof

Assume that $\models A$

Let b_1, b_2, \dots, b_n be all propositional variables that occur in the formula A , i.e.

$$A = A(b_1, b_2, \dots, b_n)$$

By the **Main Lemma** we know that, for **any** truth assignment v , the corresponding formulas A', B_1, B_2, \dots, B_n can be found such that

$$B_1, B_2, \dots, B_n \vdash A'$$

Proof

Note that in this case $A' = A$ for any v since $\models A$

We have two cases.

1. If v is such that $v(b_n) = T$, then $B_n = b_n$ and

$$B_1, B_2, \dots, b_n \vdash A$$

2. If v is such that $v(b_n) = F$, then $B_n = \neg b_n$ and by the **Main Lemma**

$$B_1, B_2, \dots, \neg b_n \vdash A$$

So, by the **Deduction Theorem** we have

$$B_1, B_2, \dots, B_{n-1} \vdash (b_n \Rightarrow A)$$

and

$$B_1, B_2, \dots, B_{n-1} \vdash (\neg b_n \Rightarrow A)$$

Proof of Completeness Theorem

By formula **8**.

$$\vdash ((A \Rightarrow B) \Rightarrow ((\neg A \Rightarrow B) \Rightarrow B))$$

for $A = b_n$, $B = A$

and by **monotonicity** we have that

$$B_1, B_2, \dots, B_{n-1} \vdash ((b_n \Rightarrow A) \Rightarrow ((\neg b_n \Rightarrow A) \Rightarrow A))$$

Applying **Modus Ponens** twice we get that

$$B_1, B_2, \dots, B_{n-1} \vdash A$$

Similarly, $v^*(B_{n-1})$ may be **T** or **F**

Applying the **Main Lemma**, the **Deduction Theorem**, **monotonicity**, formula **8**. and **Modus Ponens** twice we can eliminate B_{n-1} just as we have eliminated B_n

After **n steps**, we finally obtain proof of **A** in S , i.e. we have that

$$\vdash A$$

Constructiveness of the Proof

Observe that our proof of the **Completeness Theorem** is a **constructive** one.

Moreover, we have used in it only **Main Lemma** and **Deduction Theorem** which both have a **constructive proofs**

We **can** hence **reconstruct** proofs in each case when we apply these theorems back to the **original axioms** of H_2

The same applies to the **proofs** in H_2 of all formulas **1. - 9.**

It means that for any A , such that $\models A$, the set V_A of all v restricted to A **provides** us a method of a **construction** of the **formal proof** of A in H_2 .

Example

Example

The proof of **Completeness Theorem** defines a **method** of efficiently combining $v \in V_A$ while **constructing** the proof of A

Let's consider the following **tautology** $A = A(a, b, c)$

$$((\neg a \Rightarrow b) \Rightarrow (\neg(\neg a \Rightarrow b) \Rightarrow c))$$

We present on the next slides **all steps** of the **Proof 1** as applied to A

Example

Given

$$A(a, b, c) = ((\neg a \Rightarrow b) \Rightarrow (\neg(\neg a \Rightarrow b) \Rightarrow c))$$

By the **Main Lemma** and the assumption that

$$\models A(a, b, c)$$

any $v \in V_A$ **defines** formulas B_a, B_b, B_c such that

$$B_a, B_b, B_c \vdash A$$

The proof is based on a method of using all $v \in V_A$ (there is 8 of them) to **define** a process of **elimination** of all hypothesis B_a, B_b, B_c to **construct** the proof of A , i.e. to prove that

$$\vdash A$$

Example

Step 1: elimination of B_c

Observe that by definition, B_c is c or $\neg c$ depending on the **choice** of $v \in V_A$

We **choose** two truth assignments $v_1 \neq v_2 \in V_A$ such that

$$v_1 \upharpoonright \{a, b\} = v_2 \upharpoonright \{a, b\} \quad \text{and} \quad v_1(c) = T, \quad v_2(c) = F$$

Case 1: $v_1(c) = T$

By definition $B_c = c$

By our choice, the assumption that $\models A$ and the **Main Lemma** applied to v_1

$$B_a, B_b, c \vdash A$$

By **Deduction Theorem** we have that

$$B_a, B_b \vdash (c \Rightarrow A)$$

Example

Case 2: $v_2(c) = F$

By definition $B_c = \neg c$

By our **choice**, assumption that $\models A$, and the **Main Lemma** applied to v_2

$$B_a, B_b, \neg c \vdash A$$

By the **Deduction Theorem** we have that

$$B_a, B_b \vdash (\neg c \Rightarrow A)$$

Example

By Lemma 8. for $A = c$, $B = A$ we have that

$$\vdash ((c \Rightarrow A) \Rightarrow ((\neg c \Rightarrow A) \Rightarrow A))$$

By monotonicity we have that

$$B_a, B_b \vdash ((c \Rightarrow A) \Rightarrow ((\neg c \Rightarrow A) \Rightarrow A))$$

Applying **Modus Ponens** twice to the above property and properties on the previous slide we get that

$$B_a, B_b \vdash A$$

We have **eliminated** B_c

Example

Step 2: elimination of B_b from $B_a, B_b \vdash A$

We **repeat** the **Step 1**

As before we have 2 cases to consider: $B_b = b$ or $B_b = \neg b$

We **choose** two truth assignments $w_1 \neq w_2 \in V_A$ such that

$$w_1 \upharpoonright \{a\} = w_2 \upharpoonright \{a\} = v_1 \upharpoonright \{a\} = v_2 \upharpoonright \{a\} \text{ and } w_1(b) = T, w_2(b) = F$$

Case 1: $w_1(b) = T$ and by definition $B_b = b$

By our choice, assumption that $\models A$ and the **Main Lemma** applied to w_1

$$B_a, b \vdash A$$

By **Deduction Theorem** we have that

$$B_a \vdash (b \Rightarrow A)$$

Example

Case 2: $w_2(b) = F$ and by definition $B_b = \neg b$

By choice, assumption that $\models A$ and the **Main Lemma** applied to

w_2

$$B_a, \neg b \vdash A$$

By the **Deduction Theorem** we have that

$$B_a \vdash (\neg b \Rightarrow A)$$

Example

By Lemma 8. for $A = b$, $B = A$ we have that

$$\vdash ((b \Rightarrow A) \Rightarrow ((\neg b \Rightarrow A) \Rightarrow A))$$

By monotonicity

$$B_a \vdash ((b \Rightarrow A) \Rightarrow ((\neg b \Rightarrow A) \Rightarrow A))$$

Applying **Modus Ponens** twice to the above property and properties from the previous slide we get that

$$B_a \vdash A$$

We have **eliminated** B_b

Example

Step 3: elimination] of B_a from $B_a \vdash A$

We **repeat** the **Step 2**

As before we have 2 cases to consider: $B_a = a$ or $B_a = \neg a$

We choose two truth assignments $g_1 \neq g_2 \in V_A$ such that

$$g_1(a) = T \quad \text{and} \quad g_2(a) = F$$

Case 1: $g_1(a) = T$, and by definition $B_a = a$

By the choice, assumption that $\models A$, and the **Main Lemma** applied to g_1

$$a \vdash A$$

By **Deduction Theorem** we have that

$$\vdash (a \Rightarrow A)$$

Example

Case 2: $g_2(a) = F$ and by definition $B_a = \neg a$

By the choice, assumption that $\models A$, and the **Main Lemma** applied to g_2

$$\neg a \vdash A$$

By the **Deduction Theorem** we have that

$$\vdash (\neg a \Rightarrow A)$$

Example

By Lemma 8. for $A = a$, $B = A$ we have that

$$\vdash ((a \Rightarrow A) \Rightarrow ((\neg a \Rightarrow A) \Rightarrow A))$$

Applying **Modus Ponens** twice to the above property and properties from previous slides we get that

$$\vdash A$$

We have **eliminated** B_a , B_b , B_c and constructed the **proof** of A in S

EXERCISES

Exercise 1

The **Lemma** listed formulas **1.** - **9.** that we said are needed for both proofs of the **Completeness Theorem**.

List formulas from the **Lemma** that are are **needed** for the **Proof 1** .

Exercise 2

The **Proof 1** was carried for the language $\mathcal{L}_{\{\Rightarrow, \neg\}}$.

Extend the **Proof 1** to the language $\mathcal{L}_{\{\Rightarrow, \cup, \neg\}}$ by **adding** all new CASES concerning the new connective \cup . LIST all new formulas needed to be added to the formulas used in the original Proof 1.