

# Chapter 6: Examples of Classical Propositional Tautologies

## Implication

**Modus Ponens** known to the Stoics (3rd century B.C)

$$\models ((A \wedge (A \Rightarrow B)) \Rightarrow B)$$

## Detachment

$$\models ((A \wedge (A \Leftrightarrow B)) \Rightarrow B)$$

$$\models ((B \wedge (A \Leftrightarrow B)) \Rightarrow A)$$

**Sufficient** Given an implication

$$(A \Rightarrow B),$$

$A$  is called a *sufficient condition* for  $B$  to hold.

**Necessary** Given an implication

$$(A \Rightarrow B),$$

$B$  is called a *necessary condition* for  $A$  to hold.

## Implication Names

**Simple**       $(A \Rightarrow B)$  is called *a simple implication*.

**Converse**       $(B \Rightarrow A)$  is called *a converse implication* to  $(A \Rightarrow B)$ .

**Opposite**       $(\neg B \Rightarrow \neg A)$  is called *an opposite implication* to  $(A \Rightarrow B)$ .

**Contrary**       $(\neg A \Rightarrow \neg B)$  is called *a contrary implication* to  $(A \Rightarrow B)$ .

## Laws of contraposition

$$\models ((A \Rightarrow B) \Leftrightarrow (\neg B \Rightarrow \neg A)),$$

$$\models ((B \Rightarrow A) \Leftrightarrow (\neg A \Rightarrow \neg B)).$$

The laws of contraposition make it possible to replace, in any deductive argument, a sentence of the form  $(A \Rightarrow B)$  by  $\neg B \Rightarrow \neg A$ , and conversely.

## Necessary and sufficient :

We read  $(A \Leftrightarrow B)$  as

*B is necessary and sufficient for A*

because of the following tautology.

$$\models ((A \Leftrightarrow B)) \Leftrightarrow ((A \Rightarrow B) \cap (B \Rightarrow A)).$$

**Hypothetical syllogism** (Stoics, 3rd century B.C.)

$$\models (((A \Rightarrow B) \cap (B \Rightarrow C)) \Rightarrow (A \Rightarrow C)),$$

$$\models ((A \Rightarrow B) \Rightarrow ((B \Rightarrow C) \Rightarrow (A \Rightarrow C))),$$

$$\models ((B \Rightarrow C) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))).$$

**Modus Tollendo Ponens** (Stoics, 3rd century B.C.)

$$\models (((A \cup B) \cap \neg A) \Rightarrow B),$$

$$\models (((A \cup B) \cap \neg B) \Rightarrow A)$$

**Duns Scotus** (12/13 century)

$$\models (\neg A \Rightarrow (A \Rightarrow B))$$

**Clavius** (16th century)

$$\models ((\neg A \Rightarrow A) \Rightarrow A)$$

**Frege** (1879, first formulation of the classical propositional logic as a formalized axiomatic system )

$$\models (((A \Rightarrow (B \Rightarrow C)) \wedge (A \Rightarrow B)) \Rightarrow (A \Rightarrow C)),$$

$$\models ((A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C)))$$

**Apagogic Proofs** : means proofs by *reductio ad absurdum*.

**Reductio ad absurdum** : to prove  $A$  to be true, we assume  $\neg A$ .

If we get a contradiction, means we have proved  $A$  to be true.

$$\models ((\neg A \Rightarrow (B \cap \neg B)) \Rightarrow A)$$

**Implication form** : we want to prove  $(A \Rightarrow B)$  by *reductio ad absurdum*. Correctness of reasoning is based on the following tautologies.

$$\models (((\neg(A \Rightarrow B) \Rightarrow (C \cap \neg C)) \Rightarrow (A \Rightarrow B)),$$